

Variable Na I Absorption toward ρ Leo : Biased Neutral Formation in the Diffuse Interstellar Medium?¹

J. T. Lauroesch² and David M. Meyer²

Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208

jtl@elvis.astro.northwestern.edu, davemeyer@northwestern.edu

ABSTRACT

We present multi-epoch KPNO Coudé Feed observations of interstellar Na I and Ca II absorption toward the bright star ρ Leo. Comparisons of the Na I profiles observed over a period of 8 years reveal significant temporal variations in the Na I column in at least one component, implying that there is “structure” at scales of order of the proper motion (~ 12 AU). Archival *HST* Goddard High Resolution Spectrograph observations of the C I fine-structure excitation in the variable component suggest that the density is $\lesssim 20 \text{ cm}^{-3}$, significantly lower than the densities inferred in past H I 21 cm and Na I studies. We suggest that the bulk of the trace neutral species are in the density peaks within an interstellar cloud. The patchy distribution of these species naturally gives rise to the large fluctuations seen on scales of 10–1,000 AU in past temporal and binary studies. This picture predicts that the scales over which fluctuations will be observed vary as a function of the ionization rate of a species.

Subject headings: ISM: clouds – ISM: structure – stars: individual (HD 91316)

1. Introduction

The sightline toward ρ Leo (HD 91316) has long been a favorite of spectroscopists, even before the interstellar origin of the so-called “stationary” lines was established. Harper

¹Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained from the data archive at the Space Telescope Science Institute. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555.

²Visiting Astronomer, Kitt Peak National Observatory (KPNO). KPNO is a part of the National Optical Astronomy Observatories (NOAO), which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under contract to the National Science Foundation (NSF).

(1914) first noticed that the Ca II K line toward ρ Leo showed the characteristic difference in velocity attributable to “stationary” lines, although the line was much wider than was typically observed with hints that there were two blended lines present. Beals (1936) showed that the Ca II lines toward ρ Leo were indeed complex, making it one of the first three sightlines (along with ϵ and ζ Ori) where multiple components were known. Subsequently, Sanford, Merrill, & Wilson (1938) showed that the Na I lines also have multiple components, while Adams (1943) showed that the “violet” Ca II component of Beals (1936) was itself multiple. The velocities and equivalent widths for the Na I and Ca II lines measured from photographic spectra by Routly & Spitzer (1952) are (to within the errors) the same as those measured today. Optical observations of ρ Leo have continued to the present day with a multitude of different instruments, and have shown the complex nature of the Na I, K I, and Ca II profiles (Hobbs 1969, 1971, 1974; Welty, Hobbs, & Kulkarni 1994; Welty, Morton, & Hobbs 1996; Welty & Hobbs 2001). In addition, observations of numerous ultraviolet absorption lines have been made by *Copernicus*, *IUE*, and *HST*/Goddard High Resolution Spectrograph (GHRS).

Over the past decade, high spatial resolution optical and radio studies of the diffuse ISM have found strong evidence of pervasive subparsec-scale variations in interstellar absorption lines (Lauroesch 2001; Crawford 2003), although the precise interpretation of these fluctuations is not yet well understood (Heiles 1997; Elmegreen 1999; Deshpande 2000; Faison & Goss 2001; Welty & Fitzpatrick 2001; Lauroesch 2001; Crawford 2003). Initially, these observations were interpreted as ubiquitous concentrations of apparently dense atomic gas ($n_H \gtrsim 10^3 \text{ cm}^{-3}$) in otherwise diffuse sight-lines. However, recent direct measurements of the densities in a few of the optically selected clouds have suggested that in general the densities are much lower than previously inferred (Lauroesch *et al.* 1998; Lauroesch, Meyer, & Blades 2000; Welty & Fitzpatrick 2001), although observations in clouds identified by tracers of dense gas (K I) suggest that at least some of these features are associated with denser structures (Pan, Federman, & Welty 2001; Crawford 2002). Thus, it has been suggested that small-scale variations in the physical parameters may be responsible for much (if not all) of the observed variation in optical studies of small-scale structure. Recently, as part of a larger survey to identify very small scale (of order 10 AU) variations in interstellar lines, evidence for temporal (proper motion induced) variations in the interstellar Na I line toward ρ Leo have been observed. In this Letter, we will discuss these new observations as well as our analysis of archival Goddard High Resolution Spectrograph (GHRS) echelle observations of this sightline, and then discuss a simple model of neutral formation in the context of these results.

2. Observations and Data Reduction

Rho Leo is a bright ($V=3.8$) B1Ib runaway star, located at a distance of about 870 pc approximately 690 pc above the galactic plane (Diplas & Savage 1994). The proper motion is 6.6 ± 0.9 mas/yr (Perryman *et al.* 1997), which corresponds to a projected motion of 1.56 ± 0.21 AU/yr. Since 1989, we have obtained multiple observations of the interstellar Na I and Ca II absorption lines toward ρ Leo using the KPNO Coudé Feed and spectrograph at resolutions of ~ 1.4 and 3 km s^{-1} (see Table 1). These datasets have been reduced using the NOAO IRAF³ data reduction packages in the same fashion as that used by Watson & Meyer (1996) and Lauroesch & Meyer (1999). The extracted continuum-fitted spectra for all observing runs are shown in Figure 1 – note in particular the variation in the Na I profile at a heliocentric velocity of 18 km s^{-1} .

Table 1: ρ Leo Observation Log

Species	Date	Resolution (km s^{-1})	S/N ratio (per pixel)
Na I	11/95	1.4	370
	11/01	3.4	200
	11/02	3.4	220
	02/03	1.4	250
Ca II	05/89	4.8	190
	01/92	3.6	400
	11/02	3.6	330
Al III ^a	12/1992	3.6	35–55
C I ^{a,b}	03/1996	3.5	70
Zn II ^a	12/1992	3.4–3.6	70–80

^a Data obtained from the *HST* archive, obtained as parts of programs 2251 and 5882 (PI Hobbs).

^b Observations of the $\lambda\lambda$ 1328Å multiplet.

In addition to the ground-based data, we have taken advantage of existing archival GHRS spectra of ρ Leo. Observations using multiple settings of the ECH-A and ECH-B gratings taken in March 1996 and December 1992 (respectively) are available from *HST* observing programs 5882 and 2251 (PI Hobbs). These datasets were reduced using the stan-

³IRAF is distributed by NOAO, which are operated by AURA, Inc. under cooperative agreement with the NSF.




Fig. 1.— Observations of the Na I D₂ (5889Å), Na I D₁ (5895Å), and Ca II K (3933Å) lines towards ρ Leo. The top panel shows our two high resolution (1.4 km s⁻¹) observations, note in particular the variation in the Na I profile near 18 km s⁻¹, as well as the somewhat more subtle variation near -1 km s⁻¹. Also shown at the bottom of the plot is the 2003 profile divided by the 1995 profile (with an offset of -0.9 applied to the relative intensity for plotting purposes). The lower panel shows our three moderate resolution observations of the Ca II K line.

dard STSDAS routines (POFFSETS, DOPOFF, and SPECALIGN) for both the WSCAN (ECH-B) and FP-SPLIT (ECH-A) datasets, and then selected transitions were continuum fitted and analyzed (Figure 2).

The column densities, b-values, and relative velocities of the various components in both the KPNO and GHRS datasets were derived by profile fitting the continuum fitted spectra using the program `fits6p` (Welty, Hobbs, & York 1991). The wavelengths and oscillator strengths were taken from Morton (1991) and Welty *et al.* (1999), with the hyperfine splitting of the Na I D lines taken into account (Welty, Hobbs, & Kulkarni 1994). The Na I component model of Welty, Hobbs, & Kulkarni (1994) was used as a starting point for fitting the neutral species, with additional input from the Ca II component model of Welty, Morton, & Hobbs (1996) used for fitting the ionized species. The column density, b-value, and velocity of the various components were allowed to be free parameters in the initial fits – in the final fits the relative velocities of the components were fixed and the variation in the b-value limited for most components.

3. The $v \sim 18 \text{ km s}^{-1}$ Cloud

As noted above, the most striking feature when comparing the Na I profiles in Figure 1 is the strong temporal variation near 18 km s^{-1} . Also significant, however, is the lack of a corresponding variation in the observed Ca II profiles, as well as the lack of convincing Na I profile variations at other velocities. We should note here that part of the reason for including ρ Leo in our survey was the apparent lack of H I 21-cm emission associated with the 18 km s^{-1} component in the survey of Hobbs (1971), although more recent surveys show some 21-cm emission at this velocity in the region around ρ Leo (Hartmann & Burton 1997). In addition to showing clear evidence for temporal variability, the $v \sim 18 \text{ km s}^{-1}$ component is also relatively isolated and so we will concentrate on this component for the rest of the discussion.

The column densities for selected species derived for the $v \sim 18 \text{ km s}^{-1}$ component are given in Table 2. The first thing to note is the large Na I differences between the years 1989 and 2003, corresponding to a decrease of almost one-third in the column. However, there are two important caveats – first, we assumed the same b-value (0.58 km s^{-1}) for the component as that found by Welty, Hobbs, & Kulkarni (1994) using their higher resolution data. Second, and perhaps more important, the published Welty, Hobbs, & Kulkarni (1994) profile shows no sign of the weak absorption redward of this component seen in our datasets. We note that this shelf of absorption is seen in the profile published by Hobbs (1971, 1974) which has a similar resolution to our datasets. This suggests that these features may have




Fig. 2.— The C I 1328 multiplet and the Zn II 2026 line toward ρ Leo, with the 2003 Na I observation shown for scale. Note the lack of excited C I* and C I** absorption – the plots are centered on the strongest transitions at 1329.1004Å and 1329.5775Å respectively.

been over-resolved in the Welty, Hobbs, & Kulkarni (1994) data and “lost” during continuum placement, perhaps altering the derived column density. In any case, as shown in Figure 1, there are significant changes in the Na I column even over the past 8 years, with a decrease of $\sim 20\%$ if we assume the Welty, Hobbs, & Kulkarni (1994) b-value. A period of 8 years corresponds to a projected proper motion at the distance to the star of ~ 12 AU. One useful comparison that can be made immediately is to compare the $N(\text{H})$ one would estimate from the observed Na I columns, to that which would be derived using the observed $N(\text{Zn II})$. The average of the 1989 and 1995 Na I columns is $27.5 \times 10^{10} \text{ cm}^{-2}$, from which one would infer an $N(\text{H})$ of $\sim 5.5 \times 10^{19} \text{ cm}^{-2}$ (Ferlet, Vidal-Madjar, & Gry 1995). Similarly, one would estimate $N(\text{H}) \sim 4.6\text{--}7.2 \times 10^{19} \text{ cm}^{-2}$ from $N(\text{Zn II})$ (assuming a Zn depletion of 0.2 to 0.4 dex (Welty *et al.* 1999)). This consistency immediately highlights a potentially important difference between this cloud and the clouds studied previously in the ultraviolet by Lauroesch *et al.* (1998) and Lauroesch, Meyer, & Blades (2000), namely the observed $N(\text{Zn II})$ (and inferred $N(\text{H})$) is approximately an order of magnitude larger for this cloud, even though the $N(\text{Na I})$ is only a factor of 2–3 larger.

Table 2: Selected Column Densities at $v=18 \text{ km s}^{-1}$ toward ρ Leo

Species	Year	Lines/Multiplet Used ($\text{\AA}$)	$N(\text{X})$ (cm^{-2})
Na I	1989 ^a	5895	30.7×10^{10}
	1995	5889, 5895	$24.3 \pm 0.5 \times 10^{10} \text{ }^b$
	2001	5889, 5895	$20.8 \pm 1.0 \times 10^{10} \text{ }^b$
	2002	5889, 5895	$20.2 \pm 1.0 \times 10^{10} \text{ }^b$
	2003	5889, 5895	$20.9 \pm 0.6 \times 10^{10} \text{ }^b$
Al III	1992	1854, 1862	$5.9 \pm 1.1 \times 10^{11}$
C I	1996	1328	$57.7 \pm 6.2 \times 10^{11}$
C I*	1996	1328	$< 8.4 \times 10^{11} \text{ }^d$
C I**	1996	1328	$< 8.6 \times 10^{11} \text{ }^d$
Cr II	1992	2056, 2062	$1.8 \pm 0.2 \times 10^{11}$
Mg I	1992	2026	$17.6 \pm 1.8 \times 10^{11}$
Zn II	1992	2026, 2062	$13.1 \pm 1.8 \times 10^{11}$

We can now use the other species listed in Table 2 to estimate the physical conditions in this cloud. We can use the observed $N(\text{C I})$ and the upper limit on $N(\text{C I}^*)$ to place limits on the pressure and density in this cloud (Jenkins & Shaya 1979) – our upper limits to $N(\text{C I}^*)$ imply for $T \sim 100\text{--}500 \text{ K}$ (consistent with the observed b-value) pressures of $\log(p/k) < 3.3\text{--}3.8$ $\text{cm}^{-3} \text{ K}$. For an ideal gas at these temperatures, such pressures correspond to densities of

$n(\text{H}) \lesssim 20 \text{ cm}^{-3}$, similar to the densities found in previous studies (Lauroesch *et al.* 1998; Lauroesch, Meyer, & Blades 2000). In addition, the fine-structure equilibrium can also be used to estimate the minimum distance between the cloud and ρ Leo itself, since photon-pumping can significantly alter the observed level populations (Jenkins & Shaya 1979; Lauroesch *et al.* 1998). The observed upper limits to $N(\text{C I}\star)$ suggest that this cloud is $\gtrsim 10$ pc away from ρ Leo. The upper limit on the density coupled with our estimates of $N(\text{H})$ above suggest a cloud size of order 1 pc, much larger than the length scale over which we are seeing Na I fluctuations (~ 12 AU) but also smaller than the estimated distance between the cloud and ρ Leo. The observed ratio of $N(\text{Cr II})/N(\text{Zn II}) \sim 1.4$ would suggest that this cloud has a “warm cloud” depletion pattern (Welty *et al.* 1999). If we now assume a typical “warm cloud” value for the depletion of Al, we would estimate $N(\text{Al II} + \text{Al III}) \sim 1.1 \times 10^{13} \text{ cm}^{-2}$ from $N(\text{Zn II})$, which would suggest that while Al II is the dominant ionization state in this cloud there is likely to be some degree of photo-ionization. We can also use the measured neutral and Zn II column densities to estimate electron densities of 0.04, 0.15, and 0.05 cm^{-3} from C I, Na I and Mg I respectively assuming $T=100$ K and a “warm cloud” depletion pattern (Péquignot & Aldrovandi 1986; Lauroesch, Meyer, & Blades 2000). If we assume the dominant source of electrons is singly ionized carbon, these values correspond to a density of $\sim 100\text{--}400 \text{ cm}^{-3}$ at $T=100$ K, significantly in excess of the upper limit derived above.

4. Biased Neutral Formation in Interstellar Gas?

The parameters derived above for the $v \sim 18 \text{ km s}^{-1}$ component toward ρ Leo appear fairly typical for a diffuse cloud. The column density of hydrogen is $\sim 6 \times 10^{19} \text{ cm}^{-2}$, and we would estimate a cloud size of ~ 1 pc along the line of sight. The density of hydrogen appears to be of order 10 cm^{-3} , with the derived electron density and observed Al III suggesting some partial ionization of hydrogen. There is one significant problem that still remains however, namely the origin of the observed fluctuations in the Na I profile over relatively short timescales. Since the photo-ionization and recombination rates for Na I are relatively low (Péquignot & Aldrovandi 1986), one would not expect to see significant variations in the column on timescales shorter than thousands of years in an interstellar cloud. If we were thus to assume the observed fluctuations set the transverse size of the cloud, the earliest observations of ρ Leo suggest that significant Na I and Ca II absorption has been detectable at this velocity for at least 65 years, setting a minimum transverse size of 100 AU for this “structure” (at the distance of the star). The result is similar to the models suggested by Heiles (1997), namely a very long, thin sheet or filament with a thickness of order one thousandth the length. However, there is an alternative explanation for this

observed variation.

Consider an isolated diffuse interstellar cloud – it will have within it density fluctuations at scales much smaller than a parsec (Elmegreen 1999; Cho, Lazarian, & Vishniac 2002). Just as the force of gravity biases galaxies to form in the peaks of the cosmic density distribution, the neutral species in interstellar clouds are biased by the recombination rates to form on the peaks of the density distribution within a cloud. It is well known that recombination rates have a strong density dependence (Péquignot & Aldrovandi 1986), indeed this formed the basis for previous suggestions that relatively small density fluctuations could give rise to the observed small scale structure (Lauroesch & Meyer 1999; Lauroesch, Meyer, & Blades 2000). Furthermore, the C I survey of Jenkins & Tripp (2001) has identified a population of high pressure components in the ISM, which could represent these density peaks. We now ask what are some of the observational consequences of such a picture?

First we can estimate a minimum apparent length scale for the size of fluctuations observable in any trace species. The scale is just set by the distance a neutral can travel before it can be ionized, which in low density gas is of order the velocity of an atom divided by the photo-ionization rate. Assuming $v \sim 0.5 \text{ km s}^{-1}$ which is a “typical” b -value for the known temporally variable components, we derive an apparent scale of order 250 AU for Na I structures. Despite our having neglected to correct for collisions, this size is of order that observed in temporal fluctuations to date, and is consistent with the observations of Na I towards binary star systems. Furthermore, note that this scale is dependent upon the photo-ionization rate for a given species, and thus (for example) for Ca II we would obtain a scale about 5 times larger. This result very simply explains why we do not observe a corresponding variation in the Ca II profile towards ρ Leo. In addition, it suggests that species with higher photo-ionization rates such as Ca I and (perhaps) C I should show larger column density variations than species such as Na I or K I. Indeed, Crawford (2002) used observations of Ca I to suggest that the variable component towards κ Velorum arises in very dense material ($n_H \gtrsim 10^3 \text{ cm}^{-3}$).

A further consequence of this picture is that only a fraction of the total gas would reside in these peaks. Thus, adjacent sightlines should show smaller changes in the columns of the dominant ions, since the size of the peaks is only a fraction of the size of a cloud the additional column density for the dominant species would be significantly reduced. Consequently, the neutral fraction within the density peaks would be higher (perhaps much higher) than what one would calculate by taking the integrated columns, suggesting that the electron densities inferred from ionization balances are too low. Furthermore, the fine-structure equilibrium of C II and Si II would also be affected. Just like the neutral species, most of the observed C II* and Si II* absorption would arise in the density peaks, but the columns of the ground-states

would reflect the large scale distribution of gas and would include the perhaps dominant contribution by the material outside the peaks. Thus, in this picture, these methods provide only an ill-defined “average” electron density within a cloud, and would explain why there is so much difficulty in deriving self-consistent electron densities in many cases (Welty *et al.* 1999). Assumptions about partial ionization of hydrogen could then not be made using the (separately) derived electron and total hydrogen densities, since these tracers will likely be probing different portions of a cloud.

A final question is whether there is a connection between the observed small scale variations in Na I and the high pressure clumps observed by Jenkins & Tripp (2001). In this picture, the fine-structure equilibrium of C I becomes a measure of some sort of “average” density of the peaks along a sightline through a cloud. To date, there have been limited observations of the C I fine structure equilibrium associated with optically selected small scale structures (Lauroesch *et al.* 1998; Lauroesch, Meyer, & Blades 2000; Welty & Fitzpatrick 2001). Including ρ Leo, the observed Na I structures in all of these sightlines appear to arise in relatively low density gas, and not in the high pressure clumps detected by Jenkins & Tripp (2001). However, as Crawford (2003) recently noted for the temporally variable components towards both HD 32040 and HD 219188, these sightlines may be sampling a mixture of low pressure gas with a fraction (perhaps 20%) of high pressure, dense material with which the temporal variation may be associated. In the picture developed above, the observed temporal and binary star variations are arising in the neutral “halo” around one or more of these high pressure clumps, implying that these clumps are a fraction of the apparent size we calculated above. Yet another possibility is that we are seeing the remnant neutrals formed in transient high pressure clumps which have already expanded into the ambient cloud. Planned re-observations of both HD 32040 and HD 219188 will likely help in beginning to resolve this question. In any case, this suggests that the density of the majority of the gas in a cloud is smaller than that inferred from the C I observations, but that some fraction resides in higher density structures within which arises the bulk of the trace neutral species.

The authors thank the referee Ian Crawford for his valuable comments on this manuscript. It is also a pleasure to acknowledge the support of the staff of KPNO, especially Daryl Willmarth, for their assistance in obtaining this data.

REFERENCES

- Adams, W. S. 1943, ApJ, 97, 105
- Beals, C. S. 1936, MNRAS, 96, 661
- Cho, J., Lazarian, A., Vishniac, E. T. 2002, ApJ, 566, L49
- Crawford, I. 2002, MNRAS, 334, L33
- Crawford, I. 2003, Ap&SS, *in press*
- Desphande, A. A. 2000, MNRAS, 387, 199
- Diplas, A. and Savage, B. D. 1994, ApJS, 93, 211
- Elmegreen, B. G. 1999, ApJ, 527, 266
- Faison, M. D., & Goss, W. M. 2001, AJ, 121, 2706
- Ferlet, R., Vidal-Madjar, A., & Gry, C. 1985, ApJ, 298, 838
- Hartmann, D. & Burton, W. B. 1997, “Atlas of Galactic Neutral Hydrogen” (Cambridge : Cambridge University Press)
- Harper, W. E. 1914, *Publications of the Dominion Observatory Ottawa*, 1, 335
- Heiles, C. 1997, ApJ, 481, 193
- Hobbs, L. M. 1969, ApJ, 157, 165
- Hobbs, L. M. 1971, ApJ, 116, 333
- Hobbs, L. M. 1974, ApJ, 191, 381
- Jenkins, E. B., & Shaya, E. J. 1979, ApJ, 231, 55
- Jenkins, E. B., & Tripp, T. M. 2001, ApJS, 137, 297
- Lauroesch, J. T., Meyer, D. M., Watson, J. K., & Blades, J. C. 1998, ApJ, 507, L89
- Lauroesch, J. T., & Meyer, D. M. 1999, ApJ, 519, L181
- Lauroesch, J. T., Meyer, D. M., & Blades, J. C. 2000, ApJ, 543, L43

- Lauroesch, J. T. 2001, in the 17th IAP Astrophysics Colloquium, Gaseous Matter in Galaxies and Intergalactic Space, ed. Roger Ferlet, Martin Lemoine, Jean-Michel Desert, & Brigitte Raban (Paris: Frontier Group), 59
- Morton, D. C. 1991, *ApJS*, 77, 119
- Pan, K., Federman, S. R., & Welty, D. E. 2001, *ApJ*, 558, L105
- Pequignot, D., & Aldrovandi, S. M. V. 1986, *A&A*, 161, 169
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., Hoeg, E., Bastian, U., Bernacca, P. L., Cr  z  , M., Donati, F., Grenon, M., van Leeuwen, F., van der Marel, H., Mignard, F., Murray, C. A., Le Poole, R. S., Schrijver, H., Turon, C., Arenou, F., Froeschl  , M., & Petersen, C. S. 1994, *A&A*, 323, L49
- Routly, P. M., & Spitzer, L. 1952, *ApJ*, 115, 227
- Sanford, R. F., Merrill, P. W., & Wilson, O. C. 1938, *PASP*, 50, 58
- Watson, J. K., & Meyer, D. M. 1996, *ApJ*, 473, L127
- Welty, D. E., Hobbs, L. M., & York, D. G. 1991, *ApJS*, 75, 425
- Welty, D. E., Hobbs, L. M., & Kulkarni, V. P. 1994, *ApJ*, 436, 152
- Welty, D. E., Morton, D. C., & Hobbs, L. M. 1996, *ApJS*, 106, 533
- Welty, D. E., Hobbs, L. M., Lauroesch, J. T., Morton, D. C., Spitzer, L., & York, D. G. 1999, *ApJS*, 124, 465
- Welty, D. E., & Fitzpatrick, E. L. 2001, *ApJ*, 551, L175
- Welty, D. E., & Hobbs, L. M. 2001, *ApJS*, 133, 345